

Wound Healing Herbal Formulation from Essential Oil of Lemongrass (*Cymbopogon citratus*)

Wisdawati Wisdawati^{1*}, Rais Razak²

¹Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Universitas Muslim Indonesia, Makassar, 90231, Indonesia

²Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Muslim Indonesia, Makassar, 90231, Indonesia

*Corresponding Author: wisdawati.wisdawati@umi.ac.id

ARTICLE INFO

Article history:

Received : 13 Desember 2023

Revised : 05 September 2024

Accepted : 14 November 2024

Available online 16 November 2024

E-ISSN: [2620-3731](https://doi.org/10.32734/idjpcr.v7i1.14759)

P-ISSN: [2615-6199](https://doi.org/10.32734/idjpcr.v7i1.14759)

How to cite:

Wisdawati, W., & Razak, R. (2024). Wound healing herbal formulation from essential oil of lemongrass (*Cymbopogon citratus*). *Indonesian Journal of Pharmaceutical and Clinical Research*, 7(1), 1-6.

ABSTRACT

Untreated wound may lead to the infection. In some countries, plants are used to cure wound traditionally. Lemongrass (*Cymbopogon citratus*) has various bioactive compounds, including citronellol, limonene and geraniol that perform some pharmacological activities. In this study, the essential oil of lemongrass was designed into ointment formulation since lemongrass has been reported to have a wound healing effect in mice. The wound healing ointment made from different concentration of the essential oil of *Cymbopogon citratus* (F1: EOCC 5%, F2: EOCC 10% and F3: EOCC 15%) using a water-removable base and the characteristic as well as the stability (texture, color, smell, homogeneity, and pH) after been stored for 14 days was also evaluated. The result showed that all formulation (F1, F2, and F3) of wound healing formulation revealed good texture, consistency, odor, color, homogeneity and compatible to the skin until 14 days of storage. However, F3 (EOCC 15%) has a strong lemongrass smell compare to the other formulations.

Keyword: wound healing, ointment, essential oil, lemongrass, *Cymbopogon citratus*

ABSTRAK

Luka yang tidak ditangani dapat menyebabkan infeksi. Pengobatan luka dengan menggunakan tanaman telah lama digunakan di beberapa negara di seluruh dunia. Sereh telah diketahui memiliki beberapa senyawa bioaktif seperti citronellol, limonene dan geraniol yang telah dibuktikan memiliki aktivitas farmakologi tertentu. Pada penelitian ini, sereh dibuat formulasi sediaan salep karena telah terbukti dari beberapa penelitian dapat mengobati luka pada mencit. Salep luka dibuat dari beberapa variasi konsentrasi dari minyak atsiri herba sereh (F1: 5%, F2: 10% dan F3: 15%) menggunakan basis mudah dicuci. Selain itu, juga dilakukan evaluasi terhadap sediaan salep yang meliputi uji organoleptic (konsistensi, warna, dan bau), uji homogenitas, dan uji pH selama 14 hari. Hasil uji evaluasi menunjukkan bahwa semua formula sediaan salep (F1, F2, dan F3) memiliki konsistensi, warna, dan bau yang memenuhi kriteria. Semua formula sediaan salep (F1, F2 dan F3) juga menunjukkan homogen serta pH 7 yang sesuai dengan pH kulit sehingga tidak mengiritasi kulit. F3 (15%) menunjukkan wangi sereh yang lebih kuat dibandingkan dengan F1 (5%) dan F2 (10%).

Keyword: obat luka, salep, minyak atsiri, sereh, *Cymbopogon citratus*



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International.

<http://doi.org/10.32734/idjpcr.v7i1.14759>

1. Introduction

Skin lesion is causing by some factors, such as skin abrasion, surgical incision, burn wound, or pathology condition like diabetes or the other vascular diseases [1]. Cutaneous wound healing is a complex physiological process that involving some kinds of cells in order to repair and regenerate the damaged tissue [2], [3]. According to Departemen Kesehatan Republic Indonesia (Indonesian Health Department), the prevalence of skin injured in Indonesia in 2013 is around 8.2%.South

Sulawesi Selatan is the highest (12.8%), whereas the lowest in Jambi by 4.5% [4]. Wound healing process begins with hemostasis, and followed by inflammation, angiogenesis, cell proliferation, tissue regeneration and remodelling [2], [5].

Untreated wound will lead to a significant unpleasant feeling to the patient, infection and hypoxia[3], [5]. Traditionally, people in many countries cure the wound using plants. These plants have been reported to have a wound healing activity by different mechanisms, like anti-inflammation, increase angiogenesis, stimulate the collagen production, improve re-epithelialization process, alleviate the production of fibroblast, and vascularisation [2], [5], [6].

Lemongrass (*Cymbopogon citratus*) has various bioactive compounds, namely myrcene, citronellal, citronellol, caryophyllene, oxobisabolene, limonene, citral dan geraniol [7], [8], [9]. Therefore, lemongrass could be developed as a drug. Some studies reported that lemongrass extract has a wound healing activity by improving the production of fibroblast in mice [10]. However, the essential oil of *Cymbopogon citratus* (EOCC) is irritating when applying directly to the skin. Therefore, it needs to design into pharmaceutical formulation, like an ointment. In addition, there is a limited study regarding the wound healing formulation of EOCC.

2. Method

2.1 Sample preparation

To obtain the essential oil of *Cymbopogon citratus*, 750 gram of fresh lemongrass herba was extracted using hydro-distillation (Cherys XT ®) method. The obtaining oil then separated by adding the anhydrate sodium sulfat in order to get the pure lemongrass oil [11]. The total essential oil content of *Cymbopogon citratus* was determined using the following formula [12]:

$$\text{Total essential oil content (\%)} = \frac{\text{volume of oil (ml)}}{\text{weight of fresh sample before extraction (gram)}} \times 100 \quad (1)$$

2.2 Formulation of Wound Healing Ointment

The wound healing ointment was made by mixing the cetyl alcohol (Technical) with vaseline album (Technical) and melted then was added with propyl paraben (Technical). Sodium lauryl sulfate (Technical) was dissolved in water (Liquid Pharmalab Indonesia®), after that metil paraben was homogeneously mixed, three different concentration (5, 10 and 15 %) of EOCC was added. All the ingredients were used as seen in the Table 1.

Table 1. Pre-formulation of EOCC ointment

Material	F1 (%)	F2 (%)	F3 (%)
EOCC	5	10	15
Vaseline album	23	23.8	23.8
Cetyl alcohol	23	23.8	23.8
Sodium lauril sulfate	0.96	0.96	0.96
Propilene glycol	11.14	11.14	11.14
Metyl paraben	0.02	0.02	0.02
Propyl paraben	0.012	0.012	0.012
Water	Ad 100	Ad 100	Ad 100

Preformulation of EOCC ointments, F1: EOCC 5%, F2: EOCC 10%, and F3: EOCC 15%

2.3 The characteristic evaluation of EOCC ointment

2.3.1 Organoleptic Test of EOCC Ointment

To determine the good characteristic of EOCC ointment, some test like organoleptic, to reveal the smell, consistency, and the color after 14 days storage was conducted [13], [14], [15], [16].

2.3.2 pH evaluation

One g of EOCC ointment was dissolved in 10 mL water then the pH paper strip (NESCO ®) was used to evaluate the pH value of EOCC ointment [13], [14], [15], [16].

2.3.3 Homogeneity test

The EOCC ointment was spread in the object glass and observed for the lump appearance [13], [14], [15], [16].

3. Result and Discussion

Cymbopogon citratus revealed to have numerous pharmacological activities due to the chemical compounds found in its oil, such as citral, isoneral, isogeraniol, geraniol, geranyl acetate, citronellal, citronellol, germacrene-D, and elemol [17]. Additionally, lemongrass extract has the wound healing activity performed in mice [18]. In order to make ointment formulation of EOCC, the lemongrass was extracted using hydrodistillation method. The EOCC content was calculated based on equation (1), the EOCC content found 2% v/w (Table 2). According to Farmakope Herbal Indonesia, essential oil of lemongrass to use as a raw material in herbal product should not less than 0.15% v/w [12].

Table 2. EOCC content using hydrodistillation

Sample	Weight (gram)	EOCC (ml)	EOCC content (%v/w)
Lemongrass Herb	750	15	2

EOCC content after extraction using hydrodistillation

Since the ointment has some benefits, nonreactive and compatible with a wide range of drugs, aesthetically appealing, easy to apply and non-greasy, Remains in contact with the skin until removal is desired, then is removed easily [19]. Therefore, in this study, EOCC was designed to make as an ointment. This wound healing formulation was made from different concentration of EOCC (F1: 5%, F2: 10%, and F3: 15%). All the formulations was made 25 gram, and made triplicate. In this study, we evaluated the characteristic of wound healing ointment from EOCC, including the texture, color, odor, pH, and homogeneity, also, for stability test under room temperature for 14 days (Table 3).

Table 3. Characteristic Evaluation Result of EOCC

Formulation	Organoleptic Result						Homogeneity	
	Consistency		Color		Odor		Day 0	Day 14
	Day 0	Day 14	Day 0	Day 14	Day 0	Day 14		
F1	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
F2	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
F3	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous
	Thick and soft	Thick and soft	White	White	Lemongrass smell	Lemongrass smell	Homogeneous	Homogeneous


F1: EOCC 5%, F2: EOCC 10%, F3: EOCC 15%

The wound healing ointment used vaseline (white petrolatum) as a base, that may improve and keep the moisturizing of the skin [19]. According to Susanti, ointment base using vaseline has a good characteristic [13]. Furthermore, this formulation is a water removable base that has some advantages, easy to remove from the skin by water, non-greasy and thus provide a pleasant appearance [19], [20]. Vaseline is recommended in formulations due to non-irritating effect whereas cetyl alcohol is an emollient that maintain the moisture of the skin and thus will lead to make the skin feeling soft and smooth. The combination of Vaseline and cetyl alcohol will provide a good texture and consistency in ointment formulation. Align with this, the result of this study as seen on table 3, all the formulations (F1, F2 and F3) performed a good texture (thick and soft). Moreover, they are also non-greasy when applied to the skin. In addition, all the formulations have a white color (not dark) and thus gives the compelling appearance since this white color is one of the advantage of white petrolatum (Vaseline)[19].

EOCC has a good smell even in an ointment base. Hence, the EOCC ointment provided a pleasant to be used. Several essential oils contribute to this interesting smell, including geraniol, citronellal, citral, geranyl acetate [17], [21]. All the formulations have lemongrass smell and it appears by dose-dependent manner and no change after 14 days.

The homogeneity of the EOCC ointment was evaluated by applying the ointment in an object glass, the small particles in the mixed ingredients were investigated [13], [14], [15], [16]. Based on the Table 3, no lumps were found and indicated that all the ingredients were mixed well. Therefore, this formulation has a good homogeneity. According to figure 1, pH value of ointment formulations (F1, F2 and F3) shows the ointment compatible in the skin, as required in skin products [22], [23].

For the stability test, to investigate the stability under normal condition (room temperature) chemically and physically, the wound healing ointments (F1, F2 and F3) have been stored for 14 days. Based on result, presented in table 3 and figure 1, all the ointments has a good stability, no change in color, texture, homogeneity and pH.



F1: EOCC 5%, F2: EOCC 10%, F3: EOCC 15%, a). pH evaluation in day 0 (before being stored), b). pH evaluation in day 14 (after being stored)

4. Conclusion

All the formulations (F1, F2 and F3) of wound healing ointment from EOCC revealed good criteria of physical appearance, including texture, color, smell, also homogeneous and compatible to the skin (pH 7). Moreover, all the formulations showed the same characteristic after evaluated in 14 days.

Acknowledgements

This research has been funded by LP2S (Lembaga Pengembangan dan Sumberdaya) Universitas Muslim Indonesia (UMI).

Conflict of Interest

The authors declare that no conflict of interest for this study.

References

- [1] E. M. Tottoli, R. Dorati, I. Genta, E. Chiesa, S. Pisani, and B. Conti, 'Skin Wound Healing Process and New Emerging Technologies for Skin Wound Care and Regeneration', *Pharmaceutics*, vol. 12, no. 8, p. 735, Aug. 2020, doi: 10.3390/pharmaceutics12080735.

- [2] A. C. de O. Gonzalez, T. F. Costa, Z. de A. Andrade, and A. R. A. P. Medrado, 'Wound healing-A literature review', *An. Bras. Dermatol.*, vol. 91, pp. 614–620, 2016.
- [3] G. Han and R. Ceilley, 'Chronic Wound Healing: A Review of Current Management and Treatments', *Adv. Ther.*, vol. 34, no. 3, pp. 599–610, 2017, doi: 10.1007/s12325-017-0478-y.
- [4] R. Wintoko and A. D. N. Yadika, 'Manajemen terkini perawatan luka', *J. Kedokt. Univ. Lampung*, vol. 4, no. 2, pp. 183–189, 2020.
- [5] A. Shedoeva, D. Leavesley, Z. Upton, and C. Fan, 'Wound Healing and the Use of Medicinal Plants', *Evid. Based Complement. Alternat. Med.*, vol. 2019, p. 2684108, 2019, doi: 10.1155/2019/2684108.
- [6] V. E. Kaban, N. Nasri, Z. Rani, N. Suci, E. S. K. Sekali, and H. U. B. Sagala, 'The effect of turmeric parent extract gel (*Curcuma longa* Linn) on incision wound healing in male white rats (*Rattus norvegicus*)', *J. Pharm. Sci.*, pp. 616–627, 2024.
- [7] O. S. Oladeji, F. E. Adelowo, D. T. Ayodele, and K. A. Odelade, 'Phytochemistry and pharmacological activities of *Cymbopogon citratus*: A review', *Sci. Afr.*, vol. 6, p. e00137, Nov. 2019, doi: 10.1016/j.sciaf.2019.e00137.
- [8] E. Lulekal *et al.*, 'Phytochemical analysis and evaluation of skin irritation, acute and sub-acute toxicity of *Cymbopogon citratus* essential oil in mice and rabbits', *Toxicol. Rep.*, vol. 6, pp. 1289–1294, 2019, doi: 10.1016/j.toxrep.2019.11.002.
- [9] K. Manvitha and B. Bidya, 'Review on pharmacological activity of *Cymbopogon citratus*', *Int. J. Herb. Med.*, vol. 1, pp. 5–7, 2014.
- [10] D. S. F. Paqita, 'Pengaruh Ekstrak Sereh (*Cymbopogon Citratus*) terhadap Jumlah Fibroblas pada Penyembuhan Luka Mukosa Oral Tikus Putih', Universitas Sumatera Utara, 2021.
- [11] P. A. Handayani and H. Nurcahyanti, 'Ekstraksi minyak atsiri daun zodia (*Evodia suaveolens*) dengan metode maserasi dan distilasi air', *J. Bahan Alam Terbarukan*, vol. 3, no. 1, pp. 1–7, 2014.
- [12] Kementerian Kesehatan Republik Indonesia, *Farmakope Herbal Indonesia*, 2nd ed. Jakarta, 2017.
- [13] S. Susanti, W. Hajrin, and N. I. Hanifa, 'FORMULASI DAN EVALUASI SEDIAAN SALEP EKSTRAK ETANOLIK DAUN TEKELAN (*Chromolaena odorata* L.)', *J. Ilmu Farm. Dan Farm. Klin.*, vol. 19, no. 2, pp. 88–94, 2022.
- [14] D. P. Astuti, P. Husni, and K. Hartono, 'Formulasi dan uji stabilitas fisik sediaan gel antiseptik tangan minyak atsiri bunga lavender (*Lavandula angustifolia* Miller)', *Farmaka*, vol. 15, no. 1, pp. 176–184, 2017.
- [15] N. Lumentut, H. J. Edi, and E. M. Rumondor, 'Formulasi dan Uji Stabilitas Fisik Sediaan Krim Ekstrak Etanol Kulit Buah Pisang Goroho (*Musa acuminata* L.) Konsentrasi 12.5% Sebagai Tabir Surya', *J. MIPA*, vol. 9, no. 2, p. 42, Mar. 2020, doi: 10.35799/jmuo.9.2.2020.28248.
- [16] D. Maliana, N. Nuryanti, and H. Harwoko, 'Formulasi sediaan krim antioksidan ekstrak etanolik daun Alpukat (*Persea americana* Mill.)', *Acta Pharm. Indones.*, vol. 4, no. 2, pp. 7–15, 2016.

- [17] M. Mukarram *et al.*, ‘Lemongrass Essential Oil Components with Antimicrobial and Anticancer Activities’, *Antioxidants*, vol. 11, no. 1, p. 20, Dec. 2021, doi: 10.3390/antiox11010020.
- [18] M. Hairi, N. Dewi, and H. Khatimah, ‘Pengaruh ekstrak sereh (*Cymbopogon citratus*) terhadap panjang luka mukosa labial mencit secara klinis’, *Dentino J. Kedokt. Gigi*, vol. 1, no. 2, pp. 90–95, 2016.
- [19] M. De Villiers, ‘Ointment Bases’, 2009, pp. 277–290.
- [20] V. E. Kaban, N. Nasri, K. Gurning, H. D. Syahputra, and Z. Rani, ‘Formulasi Sediaan Lip Cream Ekstrak Daun Miana (*Coleus scutellarioides* [L] Benth.) sebagai Pewarna Alami’, *INSOLOGI J. Sains Dan Teknol.*, vol. 1, no. 4, pp. 393–400, 2022.
- [21] T. T. K. Ngan, T. T. Hien, P. H. Danh, L. T. H. Nhan, and L. X. Tien, ‘Formulation of the Lemongrass (*Cymbopogon citratus*) essential oil-based eco-friendly diffuse solution’, in *IOP Conference Series: Materials Science and Engineering*, IOP Publishing, 2020, p. 012024.
- [22] S. I. P. Rahmani and A. K. Zulkarnain, ‘Optimization of HPMC and Na-CMC as Gelling Agents on Physical Properties and Stability in Sunflower Seed Oil Gel Formulation’, *J. Food Pharm. Sci.*, vol. 11, no. 2, Jun. 2023, doi: 10.22146/jfps.8227.
- [23] R. Fitri, H. D. Syahputra, N. Nasri, V. E. Kaban, and Z. Rani, ‘Formulation of a biocellulose mask containing the essence of Aloe vera (L.) Burm. f combination with vitamin E as anti-aging’, *Sci. Pharm. Sci.*, no. 6 (40), pp. 36–42, 2022.